



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Distinct haplotype structure at the innate immune receptor Toll-like receptor 2 (TLR2) across bank vole populations and lineages in Europe

Morger, Jennifer ; Raberg, L ; Hille, S M ; Helsen, S ; Štefka, J ; Al-Sabi, M M ; Kapel, C M O ; Mappes, T ; Essbauer, S ; Ulrich, R G ; Bartolommei, P ; Mortelliti, A ; Balčiauskas, L ; van den Brink, N W ; Rémy, A ; Bajer, A ; Cheprakov, M ; Korva, M ; García-Pérez, A L ; Biek, R ; Withenshaw, S ; Tschirren, Barbara

Abstract: Parasite-mediated selection may contribute to the maintenance of genetic variation at host immune genes over long time scales. To date, the best evidence for the long-term maintenance of immunogenetic variation in natural populations comes from studies on the major histocompatibility complex (MHC) genes, whereas evidence for such processes from other immune genes remains scarce. In the present study, we show that, despite pronounced population differentiation and the occurrence of numerous private alleles within populations, the innate immune gene Toll-like receptor 2 (TLR2) displays a distinct haplotype structure in 21 bank vole (*Myodes glareolus*) populations across Europe. Haplotypes from all populations grouped in four clearly differentiated clusters, with the three main clusters co-occurring in at least three previously described mitochondrial lineages. This pattern indicates that the distinct TLR2 haplotype structure may precede the split of the mitochondrial lineages 0.19–0.56 Mya and suggests that haplotype clusters at this innate immune receptor are maintained over prolonged time in wild bank vole populations.

DOI: <https://doi.org/10.1111/bij.12593>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-117252>

Journal Article

Published Version

Originally published at:

Morger, Jennifer; Raberg, L; Hille, S M; Helsen, S; Štefka, J; Al-Sabi, M M; Kapel, C M O; Mappes, T; Essbauer, S; Ulrich, R G; Bartolommei, P; Mortelliti, A; Balčiauskas, L; van den Brink, N W; Rémy, A; Bajer, A; Cheprakov, M; Korva, M; García-Pérez, A L; Biek, R; Withenshaw, S; Tschirren, Barbara (2015). Distinct haplotype structure at the innate immune receptor Toll-like receptor 2 (TLR2) across bank vole populations and lineages in Europe. *Biological Journal of the Linnean Society*, 116(1):124-133. DOI: <https://doi.org/10.1111/bij.12593>

1 **Distinct haplotype structure at the innate immune receptor Toll-like**
2 **receptor 2 (*TLR2*) across bank vole populations and lineages in Europe**

3

4 Jennifer Morger¹, Lars Råberg², Sabine M. Hille³, Sanne Helsen⁴, Jan Štefka⁵,
5 Mohammad M. Al-Sabi⁶, Christian M. O. Kapel⁶, Tapio Mappes⁷, Sandra
6 Essbauer⁸, Rainer G. Ulrich⁹, Paola Bartolommei¹⁰, Alessio Mortelliti^{11,12},
7 Linas Balčiauskas¹³, Nico W. van den Brink^{14,15}, Alice Rémy¹⁶, Anna Bajer¹⁷,
8 Mihail Cheprakov¹⁸, Misa Korva¹⁹, Ana L. García-Pérez²⁰, Roman Biek²¹,
9 Susan Withenshaw²² & Barbara Tschirren¹

10

11 ¹ Institute of Evolutionary Biology and Environmental Studies, University of
12 Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

13 ² Department of Biology, Lund University, Sölvegatan 35, 223 62 Lund,
14 Sweden

15 ³ Institute of Wildlife Biology and Game Management, University of Natural
16 Resources and Life Sciences, Gregor Mendel-Strasse 33, 1180 Vienna,
17 Austria

18 ⁴ Evolutionary Ecology Group, Department of Biology, University of Antwerp,
19 Groenenborgerlaan 171, 2020 Antwerp, Belgium

20 ⁵ Biology Centre ASCR, Institute of Parasitology and University of South
21 Bohemia, Faculty of Science, Branišovská 31, 37005, České Budějovice,
22 Czech Republic

23 ⁶ Department of Plant and Environmental Sciences, Faculty of Science,
24 University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C,
25 Denmark

26 ⁷ Department of Biological and Environmental Science, University of
27 Jyväskylä, PO Box 35, 40014 Jyväskylä, Finland

28 ⁸ Bundeswehr Institute of Microbiology, Department Virology and
29 Rickettsiology, Neuherbergstr. 11, 80937 Munich, Germany

30 ⁹ Friedrich-Loeffler-Institut, Federal Research Institute for Animal Health,
31 Institute for Novel and Emerging Infectious Diseases, Südufer 10, 17493
32 Greifswald - Insel Riems, Germany

33 ¹⁰ Fondazione Ethoikos, Convento dell'Osservanza, 53030 Radicondoli,
34 Siena, Italy

35 ¹¹ Department of Biology and Biotechnology "Charles Darwin", University of
36 Rome "La Sapienza", Viale dell'Università 32, 00185, Rome, Italy

37 ¹² Fenner School of Environment and Society, Australian Research Council
38 Centre for Environmental Decisions, National Environmental Research
39 Program, The Australian National University, Canberra, ACT 0200, Australia

40 ¹³ Nature Research Centre, Akademijos 2, 08412 Vilnius, Lithuania

41 ¹⁴ Alterra, Wageningen UR. PO-Box 47, 6700 AA Wageningen, The
42 Netherlands

43 ¹⁵ Wageningen University Wageningen UR. Sub-department of Toxicology.
44 PO-Box 8000, 6700 EA, Wageningen, The Netherlands

45 ¹⁶ Faculty of Applied Ecology and Agricultural Sciences, Hedmark University
46 College, Anne Evenstadsvei 80, 2480 Koppang, Norway

¹⁷ Department of Parasitology, Institute of Zoology, Faculty of Biology,

University of Warsaw, 1 Miecznikowa Street, 02-096, Warsaw, Poland

¹⁸ Institute of Plant and Animal Ecology, Ural Branch, Russian Academy of

Sciences, Str 8 Marta 202, Yekaterinburg, 620144, Russia

¹⁹ Institute of Microbiology and Immunology, Faculty of Medicine, Zaloška 4,

1000 Ljubljana, Slovenia

²⁰ NEIKER – Instituto Vasco de Investigación y Desarrollo Agrario,

Department of Animal Health, Berreaga 1, 48160 Derio, Bizkaia, Spain

²¹ Boyd Orr Centre for Population and Ecosystem Health, Institute of

Biodiversity, Animal Health and Comparative Medicine, College of Medical

Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8QQ,

United Kingdom

²² Department of Evolution, Ecology & Behaviour, Institute of Integrative

Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB, United

Kingdom

Correspondence: Barbara Tschirren, Institute of Evolutionary Biology and

Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-

8057 Zurich, Switzerland; email: barbara.tschirren@ieu.uzh.ch, phone: +41

44 635 47 77, Fax: +41 44 635 47 80

Running title: *TLR2* variation in European bank voles

70 Abstract

71 Parasite-mediated selection may contribute to the maintenance of genetic
72 variation at host immune genes over long time scales. To date, the best
73 evidence for the long-term maintenance of immunogenetic variation in natural
74 populations comes from studies on the major histocompatibility complex
75 (*MHC*), whereas evidence for such processes from other immune genes
76 remains scarce.

77 Here we show that despite pronounced population differentiation and the
78 occurrence of numerous private alleles within populations, the innate immune
79 gene Toll-like receptor 2 (*TLR2*) displays a distinct haplotype structure in 21
80 bank vole (*Myodes glareolus*) populations across Europe. Haplotypes from all
81 populations grouped in four clearly differentiated clusters, with the three main
82 clusters co-occurring in at least three previously described mitochondrial
83 lineages. This pattern indicates that the distinct *TLR2* haplotype structure may
84 precede the split of the mitochondrial lineages 0.19 – 0.56 Mya and suggests
85 that haplotype clusters at this innate immune receptor are maintained over
86 prolonged time in wild bank vole populations.

87

88 **Keywords:** balancing selection – diversity – immune gene evolution –
89 immunogenetics – maintenance of genetic variation – *Myodes glareolus* –
90 parasite resistance – phylogeography – rodents – Toll-like receptors

91 Introduction

92 Understanding how genetic variation is maintained in natural
93 populations is a fundamental quest in evolutionary biology. Immune genes
94 frequently display particularly high diversity, presumably as a result of
95 parasite-mediated selection (Haldane, 1949). There are different, non-
96 mutually exclusive processes by which parasites (i.e. parasitic and pathogenic
97 organisms and agents) may contribute to the maintenance of genetic variation
98 at host defence genes. Firstly, parasites may specialise in the most common
99 host genotype, leading to negative frequency-dependent selection (Takahata
100 & Nei, 1990; Woolhouse *et al.*, 2002). Secondly, heterozygous individuals
101 may have a selective advantage over homozygous individuals (i.e.
102 overdominance), for example because of enhanced parasite recognition
103 ability or reduced immunopathology (Doherty & Zinkernagel, 1975; Apanius *et*
104 *al.*, 1997; Khor *et al.*, 2007). Finally, variation in the composition of the
105 parasite community in time and / or space may lead to fluctuating selection,
106 and hence contribute to the maintenance of genetic variation in host
107 populations (Hedrick, 2002).

108 To date, the best evidence for the maintenance of immunogenetic
109 variation over long time scales in natural populations comes from studies on
110 the major histocompatibility complex (*MHC*) genes (Hedrick, 1998;
111 Bernatchez & Landry, 2003; Piertney & Oliver, 2006). In humans, a number of
112 other immune loci with signatures of long-term balancing selection have been
113 identified (Bubb *et al.*, 2006, Andrés *et al.* 2009). In other species, however,

evidence for such patterns outside the MHC complex remains scarce (but see e.g. Ferguson *et al.*, 2012).

Toll-like receptors (TLRs) are pattern-recognition receptors (PRRs) that play a key role in the recognition of intra- and extracellular pathogens. They recognise essential structures of pathogens (Pathogen-Associated Molecular Patterns; PAMPs) and initiate innate and adaptive immune responses (Medzhitov, 2001; Akira, Uematsu & Takeuchi, 2006). Toll-like receptor 2 (*TLR2*), for example, plays an important role in the recognition of lipopeptides, crucial components of the cell membranes of Gram-positive bacteria (Jin *et al.*, 2007). Genetic polymorphisms at *TLRs* have been associated with variation in infectious disease susceptibility in humans and laboratory animals (Texereau *et al.*, 2005; Netea, Wijmenga & O'Neill, 2012), and there is accumulating evidence that they play an important role in mediating host-parasite interactions and disease susceptibility in wildlife (Jackson *et al.*, 2009; Turner *et al.*, 2011; Tschirren *et al.*, 2013; Morger *et al.*, 2014; Fornůsková *et al.*, 2014).

It has previously been found that in a Swedish bank vole *Myodes glareolus* (Schreber, 1780; Rodentia, Cricetidae) population *TLR2* displays a high diversity, with haplotypes grouping in three distinct clusters (Tschirren *et al.*, 2013). Furthermore, voles carrying haplotypes of one particular cluster (*c*₂) had a reduced *Borrelia afzelii* infection rate, and an analysis of haplotype frequencies and tree topology strongly suggested that these 'protective' alleles had evolved under positive selection (Tschirren *et al.*, 2013). One interpretation of these patterns is that the high diversity at *TLR2* is the result

of parasite-mediated selection. However, the distinct *TLR2* haplotype structure could also be a result of neutral processes, such as drift in isolated populations followed by population admixture. Indeed, eight well-defined mitochondrial bank vole lineages have been described in Europe (Deffontaine *et al.*, 2005; Kotlík *et al.*, 2006; Deffontaine *et al.*, 2009), and there is evidence that in Southern Fennoscandia re-colonisation after the last glaciation (around 11,000 – 10,000 years BP; Jaarola, Tegelström & Fredga, 1999) occurred from separate glacial refugia, creating contact zones between bank vole lineages that have evolved genetic differences while geographically isolated (Hewitt 1999; Jaarola, Tegelström & Fredga, 1999). Here we sequenced *TLR2* from 21 bank vole populations across Europe, belonging to five previously described mitochondrial lineages, to test if we find evidence that the different *TLR2* clusters are associated with different bank vole lineages. If, on the other hand, the co-occurrence of the distinct *TLR2* clusters is a general feature of this innate immune gene, and observed in several lineages and populations on a large geographical scale, this would provide indirect evidence for the long-term maintenance of *TLR2* diversity.

Materials and Methods

Study species and sampling sites

The bank vole is a common rodent in Europe. Its distribution ranges from Fennoscandia and the British Isles to northern Spain, Italy, and Russia (Corbet & Harris, 1991). For our study, we caught 385 bank voles at 21

locations in 18 countries for tissue or hair collection (mean per location: 18.3 animals; range: 10 - 21; Table 1, S1). Trapping and tissue or hair sampling were performed in accordance with the regulations of the respective countries and performed under licenses provided by the local animal welfare committees (Table S1). Distances between locations ranged from 69 to 4454 km.

Toll-like receptor 2 (TLR2) and cytochrome b (cytb) genotyping

Total genomic DNA was extracted from the samples using a standard extraction method (Laird *et al.*, 1991). We sequenced a 1149 bp fragment of *TLR2* for all individuals ($N = 385$, Table 1) following previously published protocols (Tschirren *et al.*, 2012). The amplified fragment contains the functionally relevant sites involved in pathogen-recognition and TLR heterodimerization (Jin *et al.*, 2007; Figure S2), and we previously demonstrated molecular signatures of positive selection during the evolutionary history of rodents within this gene region (Tschirren, Råberg & Westerdahl, 2011). *TLR2* amplifications were performed in a total volume of 10 μ l containing 0.2 μ l JumpStart Taq DNA Polymerase (Sigma-Aldrich), 300 nM of the primers MgITLR2F: CATCCATCACCTGACCCTTC and MgITLR2R: CCAGTAGGAATCCTGCTCG and 25 ng of genomic DNA.

To assign the different populations to the previously described mitochondrial bank vole lineages (Deffontaine *et al.*, 2005; Kotlík *et al.*, 2006), we sequenced a 1101 bp fragment of the mitochondrial cytochrome b (*cytb*) gene for three to five individuals per population following Kotlík *et al.* (2006).

Amplifications were performed in a total volume of 10 µl containing 0.2 µl JumpStart Taq DNA Polymerase (Sigma-Aldrich), 300 nM of each primer and 25 ng of genomic DNA. The PCR protocol for *TLR2* amplification consisted of an initial denaturation step at 94°C for 1 min, followed by 35 cycles of denaturation at 94°C for 30 sec, annealing at 57°C for 30 sec, and extension at 72°C for 90 sec, with a final elongation step at 72°C for 10 min. For *cytb* we used 40 cycles and an annealing temperature of 52°C. PCR products were purified and sequenced in both directions on an ABI Prism 3730 capillary sequencer (Applied Biosystems) using Big Dye terminator version 3.1 chemistry (Applied Biosystems). All *TLR2* and *cytb* raw sequences were trimmed, processed and aligned in Geneious 5.6.5 (Drummond *et al.*, 2009). Putative polymorphisms identified by the program were manually verified. *TLR2* haplotypes were reconstructed for each population separately in the program PHASE 2.1 (Stephens, Smith & Donnelly, 2001) using the default settings of a thinning interval of one, 100 burn-in iterations and 100 main iterations, and allowing for recombination (see Table S3 for additional information). If a haplotype was observed only once in the dataset, we repeated the PCR and sequencing reactions to confirm the sequence and avoid overestimation of diversity due to amplification or sequencing errors. *TLR2* and *cytb* sequences were submitted to GenBank (*TLR2* accession numbers: Table S4; *cytb* accession numbers: KJ612463 - KJ612512).

Statistical analyses

We quantified genetic diversity at *TLR2* in the 21 bank vole populations by

210 calculating the number of haplotypes, the number of synonymous and non-
211 synonymous substitutions and the number of amino acid variants using the
212 program DnaSP 5.10.01 (Librado & Rozas, 2009). To estimate population
213 differentiation at *TLR2* we calculated pairwise F_{ST} (Weir & Cockerham, 1984)
214 in Arlequin 3.5.1.3 (Excoffier & Lischer, 2010). Sequential Bonferroni
215 corrections were applied to account for multiple testing. To test if patterns of
216 haplotype frequencies deviated from neutral expectations we calculated Fay &
217 Wu's H (Fay & Wu, 2000). The empirical distribution of the test statistics was
218 generated using neutral coalescent simulations in DNAsp (Librado & Rozas,
219 2009), based on the observed number of segregating sites, 20 000 replicates
220 and allowing for a recombination rate of 0.001. A *Mus musculus TLR2*
221 sequence (NM_011905.3) was used as an outgroup.

222 We then reconstructed a *TLR2* haplotype genealogy in the program TCS 1.21
223 (Clement, Posada & Crandall, 2000) by taking synonymous and non-
224 synonymous substitutions into account. We used the resulting genealogy to
225 assign haplotypes to *TLR2* haplogroups (or clusters, defined as groups of
226 haplotypes that are separated from other groups of haplotypes by at least
227 seven mutations) following Tschirren et al. (2013). The three 'intermediate'
228 haplotypes in the Italian populations were separated by more than seven
229 mutations from the other, previously described clusters (c_1 , c_2 , c_3) (see
230 Results) and were therefore treated as a forth cluster (c_4) in the analyses. We
231 tested for deviations of *TLR2* cluster frequencies from Hardy–Weinberg
232 equilibrium within populations in Arlequin 3.5.1.2. (Excoffier & Lischer, 2010).
233 To determine to which of the previously described European bank vole

234 lineages (Deffontaine *et al.*, 2005; Kotlík *et al.*, 2006) the different study
235 populations belong, we compared the *cytb* sequences with 14 randomly
236 chosen *cytb* sequences from the seven lineages obtained from GenBank
237 (Basque: EF408066, EF408062; Spanish: EF408061, EF408060; Italian:
238 AJ639664, AJ639692; Western: DQ472319, DQ472333; Basal: DQ472327,
239 DQ472344; Eastern: DQ472347, DQ472346; Carpathian: DQ472253,
240 DQ472339). We excluded the eighth lineage (Ural) due to its large genetic
241 distance to all other lineages and because preliminary analyses showed that
242 we had no representative of this lineage in our sample. A neighbour-joining
243 tree (Maximum Composite Likelihood, 1000 bootstrap iterations) with
244 Northern red-backed vole, *Myodes rutilus* (Pallas, 1779; Rodentia, Cricetidae)
245 (JF714850) as an outgroup was created in MEGA 6 (Tamura *et al.*, 2013) and
246 used to assign the *cytb* sequences to one of the seven mitochondrial
247 lineages. To further confirm the robustness of the assignment of the study
248 populations to the different lineages, we reconstructed a *cytb* genealogy in the
249 program TCS 1.21 (Clement, Posada & Crandall, 2000) by taking
250 synonymous and non-synonymous substitutions into account. Finally, we
251 used a simulated annealing approach to identify groups that are maximally
252 differentiated from each other without taking geographical constraints into
253 account in the program SAMOVA 2.0 (Dupanloup *et al.*, 2002). This lineage
254 classification was then included in an analysis of molecular variance
255 (AMOVA) in Arlequin 3.5.1.3 (Excoffier & Lischer, 2010), which partitions the
256 total *TLR2* cluster variation into covariance fractions, resulting from the
257 variance among mitochondrial lineages, among populations within lineage,

and within populations. The number of permutations for this analysis was set to 1000.

Results

Mitochondrial lineages

The different bank vole populations could be assigned to one of five previously described mitochondrial lineages (Spanish, Italian, Western, Eastern, and Carpathian (Deffontaine *et al.*, 2005; Kotlík *et al.*, 2006) based on *cytb* sequencing (Figure 1; Figures S5, S6).

Groups that were maximally genetically differentiated from each other were identified as (SAMOVA; $K = 5$): Group 1 = ESP; Group 2 = ITA1, ITA2; Group 3 = NOR, SCO, SWE, LTU; Group 4 = RUS, UKR, DEN, FIN, POL, GER2; Group 5 = AUT, SUI, ENG, SLO, BEL, NED, GER1, CZE, which is identical to the lineage assignment of the populations based on the phylogenetic tree (Figure S5).

TLR2 haplotype diversity and differentiation

Genetic diversity at *TLR2* was high across bank vole populations with 86 haplotypes and 51 amino acid variants (Table 1). We observed a large number of private alleles within populations ($N = 67$, Table 1), even among populations in close proximity, as for example the two Italian populations ITA1 and ITA2 (Figure S7). Furthermore, we observed significant *TLR2* population

differentiation (F_{ST}) for most pairwise population comparisons (Table S11). An excess of high-frequency derived haplotypes was observed in most populations, which is indicative of positive selection (Table S8).

TLR2 clusters

TLR2 haplotypes from all European bank vole populations grouped in the same, distinct clusters (c_1 , c_2 , c_3 , c_4 ; Figure 2, Tables S4, S9). Despite the high *TLR2* diversity, large number of private alleles and strong population differentiation (see above), haplotypes of the three main clusters (c_{1-3}) were found in the Western, Eastern, and Carpathian lineages, and two of the three main clusters (c_1 , c_3) were observed in the Spanish lineage (Figure 1). Only haplotypes of one previously described cluster (c_1) were present in the Italian lineage. In addition, a fourth haplotype cluster, restricted to the Italian populations, was observed (c_4 ; Figures 1, 2).

An analysis of molecular variance (AMOVA) revealed that mitochondrial lineage explained 18% of variation in *TLR2* clusters, whereas population within lineage explained 14% of variation in *TLR2* clusters. Most variation in *TLR2* clusters was observed within populations (68%) (Table 2a). In comparison, lineage explained 55% of *cytb* variation (Table S10). When restricting the AMOVA analysis to the Western, Eastern and Carpathian lineages the proportion of variation in *TLR2* explained by lineage dropped to 0.8% (Table 2b), showing that the lineage effect in the overall analysis was due to differences in the *TLR2* composition of the Italian and Spanish clades. There was no evidence that *TLR2* cluster frequencies deviated from Hardy-

Weinberg equilibrium within populations (all P = non-significant after Bonferroni correction).

Discussion

Molecular genetic analyses revealed a high *TLR2* diversity in 21 bank vole populations across Europe. Although the haplotype composition differed markedly among populations, and a large number of private alleles was observed within populations, *TLR2* haplotypes from all populations grouped in the same distinct clusters.

Haplotypes of the three main clusters were observed in the Western, Eastern and Carpathian mitochondrial lineages, and it is possible that we underestimated *TLR2* diversity in the two other lineages (i.e. the Italian lineage where c_1 and c_4 occurred, and Spanish lineage where c_1 and c_3 occurred), because fewer samples were analysed for these groups. A molecular analysis of variance confirmed that most *TLR2* cluster variation was present within populations, rather than among populations or mitochondrial lineages. Furthermore, the variance explained by lineage was due to the very high frequency of c_3 in the Spanish lineage as well as the occurrence of a fourth cluster that was restricted to the Italian lineage. When excluding the Italian and Spanish clades, the variation explained by lineage dropped to 0.8% and was no longer significant. It shows that at least among the Eastern, Western and Carpathian clades there is no association between *TLR2* clusters and mitochondrial lineages.

329 Interestingly, the finding that *TLR2* clusters are shared among lineages
330 and populations across Europe is similar to patterns observed at the bank
331 vole MHC class II (*Dqa*-exon 2) where no genetic structure was observed at
332 the continental scale and mitochondrial lineage explained <2% of *Dqa*-exon 2
333 variation (Malé *et al.*, 2012). One interesting hypothesis is that parasites might
334 impose similar selective pressures on the two immune genes, thereby
335 maintaining genetic variation in time and space.

336 Our results suggest that the distinct *TLR2* clusters may precede the
337 split of the bank vole lineages, which is estimated to have occurred during the
338 Upper Pleistocene between 0.19 and 0.56 Mya (Deffontaine *et al.*, 2009), and
339 that clusters have been maintained within lineages since. Alternatively, *TLR2*
340 clusters may have arisen several times independently by convergent evolution
341 after the last glaciation. This, however, appears unlikely given the large
342 number of populations in which all three clusters were observed. Importantly,
343 both scenarios suggest that the co-occurrence of the three main *TLR2*
344 clusters may be favoured by natural selection. Haplotypes within clusters, on
345 the other hand, do not appear to be maintained across populations and
346 lineages, as suggested by the strong population differentiation and the large
347 number of private alleles within populations. This within-cluster differentiation
348 among population is likely the result of population isolation and drift, although
349 we can currently not exclude other evolutionary forces.

350 Only three haplotypes in the southernmost populations ITA1 and ITA2
351 did not group with one of the three previously described *TLR2* clusters. It is
352 possible that these haplotypes represent a forth cluster that was lost in other

populations during glaciation, but persisted on the Italian peninsula, which remained largely free of ice (Hewitt 1999). Alternatively, the occurrence of these 'intermediate' haplotypes could be an indication for relaxed selection in southern Europe. Indeed, a strong association between *TLR2* genotype and *Borrelia afzelii* resistance has recently been observed in bank voles (Tschirren *et al.*, 2013), and whereas *Borrelia* sp. are common tick-transmitted pathogens in most parts of Europe (Lindgren & Jaenson, 2006), they are absent in central Italy where populations ITA1 and ITA2 are situated (Cimmino *et al.*, 1992). Reduced parasite-mediated selection and / or bottlenecks during post-glacial recolonization could also explain the low *TLR2* diversity observed in populations at the northern periphery of the species range (i.e. Norway and British Isles).

Interestingly, in Spain the *Borrelia* genospecies 'R57' is highly prevalent (but absent in other parts of Europe) as is the otherwise rare *TLR2* cluster c₃ (Gil *et al.*, 2005; Barandika *et al.*, 2007). Testing for interactive effects between different *Borrelia* species and *TLR2* clusters will thus be a fruitful next step to better understand why the distinct *TLR2* clusters are maintained within bank vole lineages and populations, and why some *TLR2* clusters are common in some populations, but rare in others. Also, *TLR2* is involved in the recognition of other pathogens (Lien *et al.*, 1999; Medzhitov, 2001), which constitute multiple selective forces on *TLR2* and may contribute to the observed patterns of *TLR2* diversity and differentiation across Europe.

Besides interactions between host genotype and parasite strains or species (i.e. host G x parasite G interactions), other processes could

contribute to the maintenance of *TLR2* clusters within populations. For example, overdominance appears to play a role in the maintenance of diversity at the major histocompatibility complex (MHC) (Doherty & Zinkernagel, 1975; Apanius *et al.*, 1997; Oliver, Telfer & Piertney, 2009), and has also been suggested to contribute to the maintenance of genetic variation at TIRAP, an adaptor in the TLR signalling pathway (Khor *et al.*, 2007). However, we found no evidence for an excess of *TLR2* heterozygotes in our study populations. Alternatively, negative-frequency dependent selection might underlie the maintenance of *TLR2* clusters within populations (Tanaka & Nei, 1989; Woolhouse *et al.*, 2002).

In conclusion, our study shows that *TLR2* haplotypes group in distinct clusters in bank vole populations across Europe. Because haplotypes of the main three *TLR2* clusters were found in at least three mitochondrial lineages, our results suggest that selection may maintain genetic supertypes at this innate immune receptor in time and space.

Acknowledgements

We thank Robert J. Baker from the Texas Tech University for providing Ukrainian bank vole samples, Irene Keller and three anonymous reviewers for comments on the manuscript, and the numerous helpers that assisted with bank vole trapping and DNA extraction in the different study populations. The study was financially supported by the University of Zurich Research Priority Program 'Evolution in Action: From Genomes to Ecosystems' and the Swiss National Science Foundation (PP00P3_157455)(to BT).

401 References

- 402 **Akira S, Uematsu S, Takeuchi O. 2006.** Pathogen recognition and innate
403 immunity. *Cell* **124**: 783-801.
- 404 **Andrés AM, Hubisz MJ, Indap A, Torgerson DG, Degenhardt JD, Boyko**
405 **AR, Gutenkunst RN, White TJ, Green ED, Bustamante CD, Clark**
406 **AG, Nielsen R. 2009.** Targets of balancing selection in the human
407 genome. *Molecular Biology and Evolution* **26**: 2755-2764.
- 408 **Apanius V, Penn D, Slev PR, Ruff LR, Potts WK. 1997.** The nature of
409 selection on the major histocompatibility complex. *Critical Reviews in*
410 *Immunology* **17**: 179-224.
- 411 **Barandika JF, Hurtado A, Garcia-Esteban C, Gil H, Escudero R, Barral M,**
412 **Jado I, Juste RA, Anda P, Garcia-Perez AL. 2007.** Tick-borne
413 zoonotic bacteria in wild and domestic small mammals in northern
414 Spain. *Applied and Environmental Microbiology* **73**: 6166-6171.
- 415 **Bernatchez L, Landry C. 2003.** MHC studies in nonmodel vertebrates: what
416 have we learned about natural selection in 15 years? *Journal of*
417 *Evolutionary Biology* **16**: 363-377.
- 418 **Bubb KL, Bovee D, Buckley D, Haugen E, Kibukawa M, Paddock M,**
419 **Palmieri A, Subramanian S, Zhou Y, Kaul R, Green P, Olson MV.**
420 **2006.** Scan of human genome reveals no new loci under ancient
421 balancing selection. *Genetics* **173**: 2165-2177.
- 422 **Cimmino MA, Fumarola D, Sambri V, Accardo S. 1992.** The epidemiology
423 of Lyme borreliosis in Italy. *Microbiologica* **15**: 419-424.

424 **Clement M, Posada D, Crandall K. 2000.** TCS: a computer program to
 425 estimate gene genealogies. *Molecular Ecology* **9**: 1657-1660.

426 **Corbet GB, Harris S. 1991.** *The Handbook of British Mammals*. Blackwell
 427 Scientific Publications: Oxford.

428 **Deffontaine V, Ledevin R, Fontaine MC, Quere JP, Renaud S, Libois R,**
 429 **Michaux JR. 2009.** A relict bank vole lineage highlights the
 430 biogeographic history of the Pyrenean region in Europe. *Molecular*
 431 *Ecology* **18**: 2489-2502.

432 **Deffontaine V, Libois R, Kotlík P, Sommer R, Nieberding C, Paradis E,**
 433 **Searle JB, Michaux J. 2005.** Beyond the Mediterranean peninsulas:
 434 evidence of central European glacial refugia for a temperate forest
 435 mammal species, the bank vole (*Clethrionomys glareolus*). *Molecular*
 436 *Ecology* **14**: 1727-1739.

437 **Doherty PC, Zinkernagel RM. 1975.** Enhanced immunological surveillance in
 438 mice heterozygous at H-2 gene complex. *Nature* **256**: 50-52.

439 **Drummond AJ, Ashton B, Cheung M, Heled J, Kearse M, Moir R, Stones-**
 440 **Havas S, Thierer T, Wilson A. 2009.** Geneious v5.6.5.: Available from
 441 <http://www.geneious.com>.

442 **Dupanloup, I., Schneider, S., Excoffier, L. 2002.** A simulated annealing
 443 approach to define the genetic structure of populations. *Molecular*
 444 *Ecology* **11**: 2571-2581.

445 **Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: A new series of
 446 programs to perform population genetics analyses under Linux and
 447 Windows. *Molecular Ecology Resources* **10**: 564-567.

448 **Fay JC, Wu CI. 2000** Hitchhiking under positive Darwinian selection. *Genetics*
449 **155**, 1405–1413.

450 **Ferguson W, Dvora S, Fikes RW, Stone AC, Boissinot S. 2012.** Long-term
451 balancing selection at the antiviral gene OAS1 in Central African
452 chimpanzees. *Molecular Biology and Evolution* **29**: 1093-1103.

453 **Fornůšková A, Bryja J, Vinkler M, Macholán M, Piálek J. 2014.** Contrasting
454 patterns of polymorphism and selection in bacterial-sensing toll-like
455 receptor 4 in two house mouse subspecies. *Ecology and Evolution* **4**:
456 2931-2944.

457 **Gil H, Barral M, Escudero R, Garcia-Perez AL, Anda P. 2005.** Identification
458 of a new *Borrelia* species among small mammals in areas of northern
459 Spain where Lyme disease is endemic. *Applied and Environmental*
460 *Microbiology* **71**: 1336-1345.

461 **Guan Y, Ranoa DRE, Jiang S, Mutha SK, Li XY, Baudry J, Tapping RI.**
462 **2010.** Human TLRs 10 and 1 share common mechanisms of innate
463 immune sensing but not signaling. *Journal of Immunology* **184**: 5094-
464 5103.

465 **Haldane JBS. 1949.** Disease and evolution. *La Ricerca Scientifica* **19**: 68-76.

466 **Hedrick PW. 1998.** Balancing selection and MHC. *Genetica* **104**: 207-214.

467 **Hedrick PW. 2002.** Pathogen resistance and genetic variation at MHC loci.
468 *Evolution* **56**: 1902-1908.

469 **Hewitt GM. 1999.** Post-glacial re-colonization of European biota. *Biological*
470 *Journal of the Linnean Society* **68**: 87-112.

471 **Jaarola M, Tegelström H, Fredga K. 1999.** Colonization history in
 472 Fennoscandian rodents. *Biological Journal of the Linnean Society* **68**:
 473 113-127.

474 **Jackson JA, Friberg IM, Bolch L, Lowe A, Ralli C, Harris PD, Behnke JM,**
 475 **Bradley JE. 2009.** Immunomodulatory parasites and toll-like receptor-
 476 mediated tumour necrosis factor alpha responsiveness in wild
 477 mammals. *BMC Biology* **7**: doi:10.1186/1741-7007-1187-1116.

478 **Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik SG, Lee HY, Lee JO.**
 479 **2007.** Crystal structure of the TLR1-TLR2 heterodimer induced by
 480 binding of a tri-acylated lipopeptide. *Cell* **130**: 1071-1082.

481 **Kang JY, Nan X, Jin MS, Youn SJ, Ryu YH, Mah S, Han SH, Lee H, Paik**
 482 **SG, Lee JO. 2009.** Recognition of lipopeptide patterns by Toll-like
 483 receptor 2-Toll-like receptor 6 heterodimer. *Immunity* **31**: 873-884.

484 **Khor CC, Chapman SJ, Vannberg FO, Dunne A, Murphy C, Ling EY,**
 485 **Frodsham AJ, Walley AJ, Kyrieleis O, Khan A, Aucan C, Segal S,**
 486 **Moore CE, Knox K, Campbell SJ, Lienhardt C, Scott A, Aaby P,**
 487 **Sow OY, Grignani RT, Sillah J, Sirugo G, Peshu N, Williams TN,**
 488 **Maitland K, Davies RJO, Kwiatkowski DP, Day NP, Yala D, Crook**
 489 **DW, Marsh K, Berkley JA, O'Neill LAJ, Hill AVS. 2007.** A Mal
 490 functional variant is associated with protection against invasive
 491 pneumococcal disease, bacteremia, malaria and tuberculosis. *Nature*
 492 *Genetics* **39**: 523-528.

493 **Kotlík P, Deffontaine V, Mascheretti S, Zima J, Michaux JR, Searle JB.**
 494 **2006.** A northern glacial refugium for bank voles (*Clethrionomys*

495 *glareolus*). *Proceedings of the National Academy of Sciences of the*
496 *USA* **103**: 14860-14864.

497 **Laird PW, Zijderveld A, Linders K, Rudnicki MA, Jaenisch R, Berns A.**
498 **1991.** Simplified mammalian DNA isolation procedure. *Nucleic Acids*
499 *Research* **19**: 4293.

500 **Librado P, Rozas J. 2009.** DnaSP v5: A software for comprehensive analysis
501 of DNA polymorphism data. *Bioinformatics* **25**: 1451-1452.

502 **Lien E, Sellati TJ, Yoshimura A, Flo TH, Rawadi G, Finberg RW, Carroll**
503 **JD, Espevik T, Ingalls RR, Radolf JD, Golenbock DT. 1999.** Toll-like
504 receptor 2 functions as a pattern recognition receptor for diverse
505 bacterial products. *Journal of Biological Chemistry* **274**: 33419-33425.

506 **Lindgren E, Jaenson TGT. 2006.** Lyme borreliosis in Europe: influences of
507 climate and climate change, epidemiology, ecology and adaptation
508 measures. In: Europe WRO, ed. Copenhagen: World Health
509 Organisation. 1-35.

510 **Malé PJG, Martin JF, Galan M, Deffontaine V, Bryja J, Cosson JF,**
511 **Michaux J, Charbonnel N. 2012.** Discongruence of *Mhc* and
512 cytochrome *b* phylogeographical patterns in *Myodes glareolus*
513 (Rodentia: Cricetidae). *Biological Journal of the Linnean Society* **105**:
514 881-899.

515 **Medzhitov R. 2001.** Toll-like receptors and innate immunity. *Nature Reviews*
516 *Immunology* **1**: 135-145.

517 **Morger J, Bajnok J, Boyce K, Craig PS, Rogan MT, Lun ZR, Hide G,**
518 **Tschirren B. 2014.** Naturally occurring Toll-like receptor 11 (*TLR11*)

519 and Toll-like receptor 12 (*TLR12*) polymorphisms are not associated
520 with *Toxoplasma gondii* infection in wild wood mice. *Infection Genetics*
521 *and Evolution* **26**: 180-184.

522 **Netea MG, Wijmenga C, O'Neill LAJ. 2012.** Genetic variation in Toll-like
523 receptors and disease susceptibility. *Nature Immunology* **13**: 535-542.

524 **Oliver MK, Telfer S, Piertney SB. 2009.** Major histocompatibility complex
525 (MHC) heterozygote superiority to natural multi-parasite infections in
526 the water vole (*Arvicola terrestris*). *Proceedings of the Royal Society B*
527 **276**: 1119-1128.

528 **Piertney SB, Oliver MK. 2006.** The evolutionary ecology of the major
529 histocompatibility complex. *Heredity* **96**: 7-21.

530 **Stephens M, Smith NJ, Donnelly P. 2001.** A new statistical method for
531 haplotype reconstruction from population data. *American Journal of*
532 *Human Genetics* **68**: 978-989.

533 **Takahata N, Nei M. 1990.** Allelic genealogy under overdominant and
534 frequency-dependent selection and polymorphism of Major
535 histocompatibility complex loci *Genetics* **124**: 967-978.

536 **Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S. 2013.** MEGA6:
537 Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular*
538 *Biology and Evolution* **30**: 2725-2729.

539 **Tanaka T, Nei M. 1989.** Positive Darwinian selection observed at the
540 variable-region genes of immunoglobulins *Molecular Biology and*
541 *Evolution* **6**: 447-459.

542 **Texereau J, Chiche JD, Taylor W, Choukroun G, Comba B, Mira JP. 2005.**
 543 The importance of Toll-like receptor 2 polymorphisms in severe
 544 infections. *Clinical Infectious Diseases* **41**: S408-S415.

545 **Tschirren B, Andersson A, Scherman K, Westerdahl H, Mittl P, Råberg L.**
 546 **2013.** Polymorphisms at the innate immune receptor *TLR2* are
 547 associated with *Borrelia* infection in a wild rodent population
 548 *Proceedings of the Royal Society B* **280**: 10.1098/rspb.2013.0364.

549 **Tschirren B, Andersson M, Scherman K, Westerdahl H, Råberg L. 2012.**
 550 Contrasting patterns of diversity and population differentiation at the
 551 innate immunity gene Toll-like receptor 2 (*TLR2*) across populations of
 552 two sympatric rodent species. *Evolution* **66**: 720-731.

553 **Tschirren B, Råberg L, Westerdahl H. 2011.** Signatures of selection acting
 554 on the innate immunity gene Toll-like receptor 2 (*TLR2*) during the
 555 evolutionary history of rodents. *Journal of Evolutionary Biology* **24**:
 556 1232-1240.

557 **Turner AK, Begon M, Jackson JA, Bradley JE, Paterson S. 2011.** Genetic
 558 diversity in cytokines associated with immune variation and resistance
 559 to multiple pathogens in a natural rodent population. *PLoS Genetics* **7**:
 560 e1002343. doi:1002310.1001371/journal.pgen.1002343.

561 **Weir BS, Cockerham CC. 1984.** Estimating F-statistic for the analysis of
 562 population-structure. *Evolution* **38**: 1358-1370.

563 **Woolhouse MEJ, Webster JP, Domingo E, Charlesworth B, Levin BR.**
 564 **2002.** Biological and biomedical implications of the co-evolution of
 565 pathogens and their hosts. *Nature Genetics* **32**: 569-577.

Figure Legends

Figure 1. *TLR2* cluster frequencies and mitochondrial lineage of the sampled bank vole populations. Population IDs are given within the corresponding *TLR2* cluster pie. N = 385 animals; N = 20 – 42 alleles per population.



